

General Sobolev Inequality on Riemannian Manifold *

Qihua Ruan¹

Zhihua Chen²

1. Department of Applied Mathematics, Tongji University, Shanghai, 200092 P.R.China;
Department of Mathematics, Putian College, Putian, Fujian 351100 P.R.China.

Email: ruanqihua@163.com

2. Department of Applied Mathematics, Tongji University, Shanghai, 200092 P.R.China.

Email: zzzhzc@tongji.edu.cn

Abstract

Let M be a complete n -dimensional Riemannian manifold, if the sobolev inequalities hold on M , then the geodesic ball has maximal volume growth; if the Ricci curvature of M is nonnegative, and one of the general Sobolev inequalities holds on M , then M is diffeomorphic to R^n .

Keywords: General Sobolev inequality, Ricci curvature, diffeomorphic.

Mathematics Subject Classification (2000): 53C20, 53C21.

1. Introduction

Let M be any complete n -dimensional ($n \geq 2$) Riemannian manifold with nonnegative Ricci curvature, $C_0^\infty(M)$ be the space of smooth functions with compact support in M . Denote by dv and ∇ the Riemannian volume element and the gradient operator of M , respectively.

It is well known in [1] that Ledoux showed that: If one of the following Sobolev inequalities is satisfied,

$$(1) \quad \|f\|_p \leq C_0 \|\nabla f\|_q, \forall f \in C_0^\infty(M), \quad 1 \leq q < n, \quad \frac{1}{p} = \frac{1}{q} - \frac{1}{n}.$$

where C_0 is the optimal constant in R^n , denote $\|f\|_p$ by the L^p norm of function f ; then M is isometric to R^n .

* Project partially supported by NFSC No. 10271089; and partially supported by Fund of Department of education of Fujian Province ; No.JA04266.

The basic idea of Ledoux's result is to find a function in $C_0^\infty(M)$, then one can substitute it to (1) and obtain that $V(x_0, r) \geq V_0(r)$, here $V(x_0, r)$ denote the volume of the geodesic ball $B(x_0, r)$ of radius r with center x_0 , and $V_0(r)$ the volume of the Euclidean ball of radius r in R^n . Since the Ricci curvature of M is nonnegative, from Bishop's comparison theorem[2], we know that $V(x_0, r) \leq V_0(r)$, so M is isometric to R^n .

Later Xia combined Ledoux's method with Cheeger and Colding's result[3], which is that given an integer $n \geq 2$, there exists a constant $\delta(n) > 0$ such that any n -dimensional complete Riemannian manifold with nonnegative Ricci curvature and $V(x_0, r) \geq (1 - \delta(n))V_0(r)$ for some $x_0 \in M$ and all $r > 0$ is diffeomorphic to R^n ; and proved the following theorem [4]:

Theorem 1.1 *Let M be a complete n -dimensional Riemannian manifold with nonnegative Ricci curvature, if one of the following sobolev inequalities is satisfied,*

$$(2) \quad \|f\|_p \leq C_1 \|\nabla f\|_q, \forall f \in C_0^\infty(M), \quad 1 \leq q < n, \quad \frac{1}{p} = \frac{1}{q} - \frac{1}{n}.$$

where the positive constant $C_1 > C_0$, then M is diffeomorphic to R^n .

As we known, the Sobolev inequality(2) belongs to a general Sobolev inequalities of the type

$$(3) \quad \|f\|_t \leq C \|f\|_s^\theta \|\nabla f\|_q^{1-\theta}, \forall f \in C_0^\infty(M), \quad \frac{1}{t} = \frac{\theta}{s} + \frac{1-\theta}{p}.$$

(see [5]), where C is a positive constant. Inequality (2) corresponds to $\theta = 0$. When $q = r = 2$, and $\theta = 2/(n+2)$, it corresponds to the Nash inequality

$$(4) \quad \left(\int |f|^2 dv \right)^{1+\frac{2}{n}} \leq C \left(\int |f| dv \right)^{\frac{4}{n}} \int |\nabla f|^2 dv, \forall f \in C_0^\infty(M),$$

[see(6)], where C is a positive constant.

In [], Ledoux conjecture two problem: one is that if one of Nash inequalities is satisfied on M , then M is isometric to R^n ; the other is that without any curvature assumption, if the Sobolev inequalities are satisfied on M , then the geodesic ball has maximal volume growth, namely $\lim_{r \rightarrow +\infty} \frac{V(x_0, r)}{V_0(r)} \geq C$, where positive constant C depends only on n and q . The first problem was proved by Druet, Hebey, and Vaugon(see []). In this paper, we want to confirm the other problem, that is

Theorem 1.2 *Let M be a complete n -dimensional Riemannian manifold, if the Sobolev inequalities (2) are satisfied on M , then the geodesic ball in M has maximal volume growth.*

We also generalize Theorem 1.1 to obtain the following result:

Theorem 1.3 *Let M be a complete n -dimensional Riemannian manifold with nonnegative Ricci curvature, if one of the general Sobolev inequalities (3) is satisfied on M , then M is diffeomorphic to R^n .*

2. Proof of Theorem 1.2 and 1.3

Now we first prove Theorem 1.2.

Proof: As we known, there always exists a cut-off function $f_m \in C_0^\infty(M)$ such that $f_m \equiv 1$ on $B(x_0, 2^m r)$, $f_m \equiv 0$ outside $B(x_0, 2^{m+1} r)$, $|\nabla f_m| \leq \frac{1}{2^m r}$ on $B(x_0, 2^{m+1} r) \setminus B(x_0, 2^m r)$ for any real integer m . Substituting them to (1), we have that

$$V(x_0, 2^m r)^{\frac{1}{p}} \leq \frac{C_0}{2^m r} V(x_0, 2^{m+1} r)^{\frac{1}{q}}$$

Thus set $\frac{q}{p} = a$, we get that

$$\begin{aligned} V(x_0, r) &\geq \left(\frac{r}{2C_0}\right)^q V(x_0, \frac{r}{2})^a \\ &\geq (C_0^{-1} r)^{q(1+a)} 2^{-q(1+2a)} V(x_0, \frac{r}{2^2})^{a^2} \\ &\geq (C_0^{-1} r)^{q(1+a+a^2)} 2^{-q(1+2a+3a^2)} V(x_0, \frac{r}{2^3})^{a^3} \\ &\geq \cdots \geq (C_0^{-1} r)^{q \sum_{i=0}^k a^i} 2^{-q \sum_{i=0}^k (i+1)a^i} V(x_0, \frac{r}{2^{k+1}})^{a^{k+1}} \\ &\geq \cdots \geq (C_0^{-1} r)^{q \sum_{i=0}^k a^i} 2^{-q \sum_{i=0}^k (i+1)a^i} V(x_0, 1)^{a^{k+1}} \quad (r \geq 2^{k+1}) \end{aligned}$$

Let $k \rightarrow +\infty$, $r \rightarrow +\infty$, since $a = \frac{q}{p} < 1$, then we see that

$$\lim_{r \rightarrow +\infty} \frac{V(x_0, r)}{r^n} \geq C_0^{-n} 2^{n - \frac{n^2}{q}}$$

So we proved Theorem 1.2.

Next we prove Theorem 1.3

Proof: As proof of Theorem 1.2, there always exists a cut-off function $f \in C_0^\infty(M)$ such that $f \equiv 1$ on $B(x_0, r)$, $f \equiv 0$ outside $B(x_0, kr)$, $|\nabla f| \leq \frac{1}{(k-1)r}$ on $B(x_0, kr) \setminus B(x_0, r)$ for some $k > 1$.

Substituting it to (1), we have that

$$\begin{aligned} V(x_0, r)^{\frac{1}{t}} &\leq C V(x_0, kr)^{\frac{\theta}{s}} \left(\frac{1}{(k-1)r}\right)^{1-\theta} V(x_0, kr)^{\frac{1-\theta}{q}} \\ V(x_0, r)^{\frac{1}{t}} &\leq \frac{C}{((k-1)r)^{1-\theta}} V(x_0, kr)^{\frac{\theta}{s} + \frac{1-\theta}{q}} \\ V(x_0, r)^{\frac{1}{t}} &\leq \frac{C}{((k-1)r)^{1-\theta}} k^{n(\frac{\theta}{s} + \frac{1-\theta}{q})} V(x_0, r)^{\frac{\theta}{s} + \frac{1-\theta}{q}} \\ 1 &\leq \frac{C k^{n(\frac{\theta}{s} + \frac{1-\theta}{q})}}{((k-1)r)^{1-\theta}} V(x_0, r)^{\frac{\theta}{s} + \frac{1-\theta}{q} - \frac{1}{t}} \end{aligned}$$

$$1 \leq \frac{Ck^{n(\frac{\theta}{s} + \frac{1-\theta}{q})}}{((k-1)r)^{1-\theta}} V(x_0, r)^{\frac{1-\theta}{n}}$$

$$V(x_0, r) \geq \frac{(k-1)^n}{C^{\frac{n}{1-\theta}} k^{\frac{n^2\theta}{s(1-\theta)} + \frac{n^2}{q}}} r^n$$

For fixed number θ and q , we can choose one $k > (C^{-\frac{1}{1-\theta}} \omega_n^{-\frac{1}{n}})^{\frac{1}{\frac{n\theta}{s(1-\theta)} + \frac{n}{q} - 1}}$ such that $V(x_0, r) \geq C'V_0(r)$ and $C' < 1$. Then there always exists a $\delta(n) > 0$ such that $V(x_0, r) \geq (1 - \delta(n))V_0(r)$. Thus from Cheeger and Colding's result, we see that M is diffeomorphic to R^n .

If $\theta = 0$, the general Sobolev inequality change into the Sobolev inequality, thus we generalize Xia's result. If $q = r = 2$ and $\theta = 2/(n+2)$, then the general Sobolev inequality become the Nash inequality, so we get the following corollary.

Corollary 2.1 *Let M be a complete n -dimensional Riemannian manifold with nonnegative Ricci curvature, if one of the Nash inequalities (4) is satisfied on M , then M is diffeomorphic to R^n .*

References

- [1] M. Ledoux, On manifolds with nonnegative Ricci curvature and Sobolev inequalities, Comm. Anal. Geom. 7(1999), 347–353.
- [2] R. L. Bishop and R. J. Crittenden, Geometry of manifolds, Academic Press, New York, 1964.
- [3] J. Cheeger and T. Colding, On the structure of space with Ricci curvature bounded below I, J. Diff. Geom. 46(1997):406–480.
- [4] C. Y. Xia, Complete manifolds with nonnegative Ricci curvature and almost best Sobolev constant. Illinois J. Math. 45(2001) 1253–1259.
- [5] D. Bakry, T. Coulhon, M. Ledoux and L. Saloff-Coste, Sobolev inequalities in disguise, Indiana J. Math. 44(1995), 1033–1074.
- [6] E. B. Davies, Heat kernels and spectral theory, Cambridge Uni. Press, 1989.
- [7] R. Schoen and S. T. Yau, Lectures on differential geometry in Conference Proceedings and Lecture Notes in Geometry and Topology, Vol 1, International Press publications, 1994.
- [8] W. X. Shi, Ricci flow and the uniformization on complete noncompact *kähler* manifolds, J. Diff. Geom. 45(1997), 94–220.